Resonant Mechanics Magnetism Equations

Alignment Transport, Helicity & Chronotactic Dynamics

Zachery Claude Maarschalkerweerd Jones

Abstract

In Resonant Mechanics, magnetism is not a separate "force" but the *transport of alignment* along lane-pure, low-shear corridors of the Field. Photons carry alignment credit; helicity records signed twist; and the temporal substrate (*chronotaxy*) supplies the cadence that keeps propagation adiabatic. This report formulates a magnetism sector consistent with the core Resonant Mechanics principles: (i) *Form* (boundary code) sets acceptance and selection rules, (ii) *Field* (stored potential with helicity) provides the energetic reservoir, and (iii) *Flow* (gyroscopic–cycloidal actuation) writes coherence as motion. This report presents field definitions, equations of motion, energy and momentum balances, constitutive laws (including helicity coupling), limiting cases (reduction to Maxwell theory in vacuum), and operational roles and predictions.

Contents

1 Foundations & Notation **2**
- 1.1 Chronotaxy & Lanes . . . 2
- 1.2 Fields & Potentials . . . 2
- 1.3 Chronotactic Sources . . . 2

2 Lagrangian & Field Equations **2**
- 2.1 Magnetism Sector Lagrangian . . . 2
- 2.2 Euler–Lagrange Equations . . . 3
- 2.3 Lane-Purity Constraint . . . 3

3 Energy, Momentum & Conservation **3**
- 3.1 Energy Density & Flux . . . 3
- 3.2 Momentum Balance . . . 4

4 Constitutive Laws & Media **4**
- 4.1 Isotropic Laminar Medium . . . 4
- 4.2 Chiral / Helical Medium . . . 4
- 4.3 Coupling to Chronotactic Geometry . . . 4

5 Waves & Dispersion **5**
- 5.1 Wave Equations . . . 5
- 5.2 Vacuum Limit (Maxwell Theory) . . . 5

6 Operational Roles & Predictions **5**

7 Measurement & Inference **5**

8 Summary of the Magnetism Sector **6**

9 Conclusion **6**

1 Foundations & Notation

1.1 Chronotaxy & Lanes

- **Chronotaxy:** the local time-order field governing rephasing; we denote its proper time by τ.
- **Helicity lanes:** two single-sense channels (right/left), kept orthogonal by domain membranes to prevent lane mixing.

1.2 Fields & Potentials

We introduce *alignment potentials* $(\Phi_A, \boldsymbol{\mathcal{A}})$ and define *alignment fields*

$$\boldsymbol{\mathcal{E}} \equiv -\frac{\partial \boldsymbol{\mathcal{A}}}{\partial \tau} - \nabla \Phi_A, \qquad \boldsymbol{\mathcal{B}} \equiv \nabla \times \boldsymbol{\mathcal{A}}. \tag{1}$$

Here $\boldsymbol{\mathcal{E}}$ plays the role of an *alignment drive* and $\boldsymbol{\mathcal{B}}$ the *alignment flux* (magnetic-like). The *helicity density* is

$$\mathsf{h} \equiv \boldsymbol{\mathcal{A}} \cdot \boldsymbol{\mathcal{B}}, \tag{2}$$

which measures signed twist stored in the Field along a lane.

1.3 Chronotactic Sources

We distinguish *alignment charge* ρ_A and *alignment current* $\boldsymbol{\mathcal{J}}_A$ (phase-work sources tied to boundary acceptance):

$$\frac{\partial \rho_A}{\partial \tau} + \nabla \cdot \boldsymbol{\mathcal{J}}_A = 0 \qquad \text{(ledger continuity)}. \tag{3}$$

2 Lagrangian & Field Equations

2.1 Magnetism Sector Lagrangian

In Resonant Mechanics, the magnetism sector is a medium-like theory with helicity coupling and chronotactic time τ:

$$\mathcal{L}_{\text{mag}} = \frac{1}{2}\,\varepsilon_R\,\|\boldsymbol{\mathcal{E}}\|^2 - \frac{1}{2}\,\mu_R^{-1}\,\|\boldsymbol{\mathcal{B}}\|^2 \;+\; \eta_{\mathsf{h}}\,\mathsf{h} \;-\; \rho_A\,\Phi_A \;+\; \boldsymbol{\mathcal{J}}_A \cdot \boldsymbol{\mathcal{A}} \;+\; \lambda\,\nabla\Theta : \big(\nabla \times \boldsymbol{\mathcal{A}}\big). \tag{4}$$

Parameters:

- ε_R: alignment permittivity (chronotactic response to drive).
- μ_R: alignment permeability (storage of alignment flux).
- η_h: helicity coupling strength to h.
- Θ: chronotactic angle field; λ couples magnetism to gradients of Θ (geometric-phase feedback).

The double contraction $A : B$ denotes $\sum_{ij} A_{ij} B_{ij}$.

2.2 Euler–Lagrange Equations

Variation with respect to Φ_A and $\boldsymbol{\mathcal{A}}$ yields *Resonant–Maxwell equations*:

$$\textbf{(Gauss–align)} \quad \nabla \cdot \boldsymbol{\mathcal{D}} = \rho_A, \tag{5}$$

$$\textbf{(Faraday)} \quad \nabla \times \boldsymbol{\mathcal{E}} = -\frac{\partial \boldsymbol{\mathcal{B}}}{\partial \tau}, \tag{6}$$

$$\textbf{(No-monopole)} \quad \nabla \cdot \boldsymbol{\mathcal{B}} = 0, \tag{7}$$

$$\textbf{(Ampère–align)} \quad \nabla \times \boldsymbol{\mathcal{H}} = \frac{\partial \boldsymbol{\mathcal{D}}}{\partial \tau} + \boldsymbol{\mathcal{J}}_A + \boldsymbol{\mathcal{S}}_h, \tag{8}$$

with constitutive definitions (from (4))

$$\boldsymbol{\mathcal{D}} \equiv \frac{\partial \mathcal{L}_{\mathrm{mag}}}{\partial \boldsymbol{\mathcal{E}}} = \varepsilon_R \, \boldsymbol{\mathcal{E}}, \qquad \boldsymbol{\mathcal{H}} \equiv -\frac{\partial \mathcal{L}_{\mathrm{mag}}}{\partial \boldsymbol{\mathcal{B}}} = \mu_R^{-1} \, \boldsymbol{\mathcal{B}} - \eta_h \, \boldsymbol{\mathcal{A}}. \tag{9}$$

The helicity-induced *source* term reads

$$\boldsymbol{\mathcal{S}}_h \equiv \eta_h \nabla \times \boldsymbol{\mathcal{A}} \; + \; \nabla \times \big(\lambda \, \nabla \Theta\big), \tag{10}$$

expressing (i) direct helicity feedback and (ii) chronotactic coupling.

2.3 Lane-Purity Constraint

Lane orthogonality (no mixing of opposite helicity channels) imposes

$$\Pi_- \, \boldsymbol{\mathcal{B}} = \mathbf{0}, \qquad \Pi_- \, \boldsymbol{\mathcal{E}} = \mathbf{0}, \tag{11}$$

for the chosen lane projector Π_-. In practice, this is enforced by boundary code and membranes; numerically it acts as a constraint or penalty term.

3 Energy, Momentum & Conservation

3.1 Energy Density & Flux

The magnetism-sector energy density and flux (Poynting-like) in laminar corridors are

$$\mathcal{U}_{\mathrm{mag}} = \frac{1}{2} \, \boldsymbol{\mathcal{E}} \cdot \boldsymbol{\mathcal{D}} + \frac{1}{2} \, \boldsymbol{\mathcal{B}} \cdot \boldsymbol{\mathcal{H}}, \tag{12}$$

$$\boldsymbol{\mathcal{S}}_{\mathrm{mag}} = \boldsymbol{\mathcal{E}} \times \boldsymbol{\mathcal{H}}, \tag{13}$$

and obey the conservation law

$$\frac{\partial \mathcal{U}_{\mathrm{mag}}}{\partial \tau} + \nabla \cdot \boldsymbol{\mathcal{S}}_{\mathrm{mag}} = -\,\boldsymbol{\mathcal{J}}_A \cdot \boldsymbol{\mathcal{E}} \;-\; \boldsymbol{\mathcal{H}} \cdot \boldsymbol{\mathcal{S}}_h. \tag{14}$$

The right-hand terms represent (i) ledgered work delivered to alignment sources and (ii) helicity/chronotactic exchange.

3.2 Momentum Balance

Define the stress tensor

$$\mathbb{T}_{ij} = \mathcal{U}_{\mathrm{mag}}\,\delta_{ij} - \mathcal{D}_i \mathcal{E}_j - \mathcal{H}_i \mathcal{B}_j. \tag{15}$$

Then the momentum conservation law in the magnetism sector reads

$$\frac{\partial}{\partial\,\tau}(\boldsymbol{\mathcal{D}} \times \boldsymbol{\mathcal{B}}) + \nabla \cdot \mathbb{T} = -\,\rho_A\,\boldsymbol{\mathcal{E}} - \boldsymbol{\mathcal{J}}_A \times \boldsymbol{\mathcal{B}} - \boldsymbol{\mathcal{Q}}_h, \tag{16}$$

where $\boldsymbol{\mathcal{Q}}_h$ encodes helicity/chronotactic momentum exchange.

4 Constitutive Laws & Media

4.1 Isotropic Laminar Medium

In a homogeneous, laminar medium:

$$\boldsymbol{\mathcal{D}} = \varepsilon_R\,\boldsymbol{\mathcal{E}}, \qquad \boldsymbol{\mathcal{H}} = \mu_R^{-1}\,\boldsymbol{\mathcal{B}} - \eta_h\,\boldsymbol{\mathcal{A}}. \tag{17}$$

4.2 Chiral / Helical Medium

In a helical environment, lowest-order bianisotropy adds

$$\boldsymbol{\mathcal{D}} = \varepsilon_R\,\boldsymbol{\mathcal{E}} + \xi\,(\nabla \times \boldsymbol{\mathcal{E}}), \qquad \boldsymbol{\mathcal{H}} = \mu_R^{-1}\,\boldsymbol{\mathcal{B}} - \eta_h\,\boldsymbol{\mathcal{A}} + \zeta\,(\nabla \times \boldsymbol{\mathcal{B}}), \tag{18}$$

with small, dispersion-setting ξ, ζ (lane-pure, no mixing).

4.3 Coupling to Chronotactic Geometry

When chronotactic curvature is non-negligible, allow

$$\varepsilon_R \to \varepsilon_R(\Theta, \nabla\Theta), \qquad \mu_R^{-1} \to \mu_R^{-1}(\Theta, \nabla\Theta), \qquad \eta_h \to \eta_h(\Theta), \tag{19}$$

capturing smooth, adiabatic retunes of impedance and helicity coupling.

5 Waves & Dispersion

5.1 Wave Equations

Eliminating sources (and with constant parameters) yields

$$\frac{\partial^2 \boldsymbol{\mathcal{B}}}{\partial \tau^2} - c_R^2 \, \nabla^2 \boldsymbol{\mathcal{B}} = -\, \mu_R \, \frac{\partial \boldsymbol{\mathcal{S}}_h}{\partial \tau}, \tag{20}$$

$$\frac{\partial^2 \boldsymbol{\mathcal{E}}}{\partial \tau^2} - c_R^2 \, \nabla^2 \boldsymbol{\mathcal{E}} = -\, \frac{1}{\varepsilon_R} \, \frac{\partial \boldsymbol{\mathcal{S}}_h}{\partial \tau}, \tag{21}$$

with $c_R \equiv (\varepsilon_R \mu_R)^{-1/2}$ the lane-pure propagation speed (identifies with c in vacuum limit).

5.2 Vacuum Limit (Maxwell Theory)

For $\eta_h = \lambda = 0$, constant $\varepsilon_R = \varepsilon_0$, $\mu_R = \mu_0$, and $\tau \to t$,

$$\nabla \cdot \boldsymbol{E} = \frac{\rho}{\varepsilon_0}, \quad \nabla \times \boldsymbol{E} = -\frac{\partial \boldsymbol{B}}{\partial t}, \quad \nabla \cdot \boldsymbol{B} = 0, \quad \nabla \times \boldsymbol{B} = \mu_0 \, \boldsymbol{J} + \mu_0 \varepsilon_0 \, \frac{\partial \boldsymbol{E}}{\partial t}, \tag{22}$$

recovered as the standard Maxwell system.

6 Operational Roles & Predictions

1. **Lane-Pure Propagation:** Long-range alignment transport with negligible dispersion in healthy (laminar) corridors.
2. **Helicity Imprint:** Chiral media introduce small, controllable rotation/ellipticity tied to η_h, ξ, ζ.
3. **Chronotactic Retune:** Slow changes in Θ modulate ε_R and μ_R, shifting phase/group velocities adiabatically.
4. **No Lane Mixing:** Domain membranes enforce (11); violations diagnose damage (turbulence) rather than new physics.
5. **Poynting Closure:** The conservation law (14) provides a ledger for absorbed/emitted phase-work (photonic bookkeeping).

7 Measurement & Inference

- **Polarimetry:** Tracks lane purity and helicity coupling through rotation of alignment states.
- **Interferometry:** Measures chronotactic retunes via stable phase shifts under controlled Θ-gradients.
- **Dispersion Probing:** Broadband timing of transient alignment pulses constrains $(\varepsilon_R, \mu_R, \eta_h)$ in situ.

8 Summary of the Magnetism Sector

$$\boxed{\begin{aligned}
&\boldsymbol{\mathcal{E}} = -\frac{\partial \boldsymbol{\mathcal{A}}}{\partial \tau} - \nabla \Phi_A, \quad \boldsymbol{\mathcal{B}} = \nabla \times \boldsymbol{\mathcal{A}}, \quad \mathsf{h} = \boldsymbol{\mathcal{A}} \cdot \boldsymbol{\mathcal{B}}, \\
&\nabla \cdot \boldsymbol{\mathcal{D}} = \rho_A, \ \nabla \times \boldsymbol{\mathcal{E}} = -\frac{\partial \boldsymbol{\mathcal{B}}}{\partial \tau}, \ \nabla \cdot \boldsymbol{\mathcal{B}} = 0, \ \nabla \times \boldsymbol{\mathcal{H}} = \frac{\partial \boldsymbol{\mathcal{D}}}{\partial \tau} + \boldsymbol{\mathcal{J}}_A + \boldsymbol{\mathcal{S}}_h, \\
&\boldsymbol{\mathcal{D}} = \varepsilon_R \boldsymbol{\mathcal{E}}, \qquad \boldsymbol{\mathcal{H}} = \mu_R^{-1} \boldsymbol{\mathcal{B}} - \eta_h \boldsymbol{\mathcal{A}}, \qquad \boldsymbol{\mathcal{S}}_h = \eta_h \nabla \times \boldsymbol{\mathcal{A}} + \nabla \times (\lambda \nabla \Theta).
\end{aligned}}$$

9 Conclusion

Magnetism in Resonant Mechanics is *alignment transport* shaped by helicity, safeguarded by lane purity, and paced by chronotaxy. The sector above reduces to Maxwell theory in the vacuum limit, while extending it with helicity-ledger terms and adiabatic chronotactic retunes. Energy and momentum are conserved as counted phase-work; polarization and interferometric observables directly read out the health of corridors. In this view, magnetic phenomena are not pushes and pulls but the choreography of coherence: alignment moving cleanly through a universe that prefers to stay in tune.